

Wildlife and Habitats

Fourth Edition



Preview

Sample Essays & Tables of Contents

About the Author



Susan C. Morse, founder and science director of Keeping Track, is widely regarded as an expert in natural history and one of North America's leading wildlife trackers. For over 50 years, she has monitored wildlife, documenting the presence and habitat requirements of such elusive species as bobcat, black bear, Canada lynx, caribou, and cougar.

When she is not conducting field research, leading training programs, or photographing wildlife, Sue shares her findings and award-winning photographs with fellow professionals, conservation organizations, students and wildlife enthusiasts of all ages. A University of Vermont graduate, she has mentored numerous young people over the years, inspiring many to pursue careers in conservation.

Publications

In addition to her work with Keeping Track, Sue operates a forestry consulting business, writes natural history features, and illustrates her work with her wildlife photography. In 2026, Keeping Track published the fourth edition of *Wildlife and Habitats*, bringing together and greatly expanding upon essays Sue wrote and illustrated for *Northern Woodlands* magazine over a span of 30 years. She is also the author and primary photographer of *A Guide to Recognizing the Florida Panther, Its Tracks and Sign*.

Awards and Recognition

Sue's honors include the Environmental Leader Award from Unity College and the Franklin Fairbanks Award for her lifelong work enriching New England residents' understanding of the natural world. The Adirondack Council has also recognized Sue and Keeping Track for decades of conservation work in the Lake Champlain basin bioregion.

She and Keeping Track have been featured on National Public Radio's *Morning Edition* and in numerous publications, including *Adirondack Life*, *Audubon*, *Smithsonian*, and *Wild Earth*.

Sue lives in Jericho, Vermont, alongside a rich and diverse forest she has helped conserve. Generations of Keeping Track wildlife-monitoring teams have trained there, and students from around the world have come to learn about its ecology and wild inhabitants. The forest also serves as a living laboratory for Sue's continuing wildlife research and stewardship.



**Celebrating those who care
about wildlife!**

Sample Essays



Making Sense of Scent Marking


Story and Photographs by Susan C. Morse



What do deer, moose, cougars, bobcats, lynx and bears have in common? They communicate with others of their species through "scent-marking." In fact, nearly all terrestrial mammals from mice to elephants use "semiochemicals" so that they may olfactorily communicate with each other in absentia. Semiochemicals is from the Greek word *semeion*, meaning "signal" or "sign."

Carnivore and cervid species maintain social order through an elaborate repertoire of chemo-communications which are continually posted throughout the habitat and refreshed as needed. Urine, feces, saliva and exudates from various skin glands are deposited in specific scent marking environments which are often visually conspicuous and thereby attract olfactory inspection by neighbors and newcomers alike.

Scent marks, posted across a solitary wide-ranging carnivore's expansive home range area, makes possible necessary communication in an energetically efficient manner. Through scent marking, bobcats, cougars, or bears can exploit the resources of huge territories yet remain in near constant contact with others of their kind. Scent messages distributed throughout the shared environment mediate competition and aggressive encounters, while enabling breeding partners to locate one another at other times.



- (A) Making a scrape by scuffing with the hind feet
- (B) Facial marking with scent from submandibular glands located beneath the chin.
- (C) Female inspects urine posted by tom seeking a date.
- (D) Remote camera picture of a tom bobcat at Wolfrun facial marking a conspicuous portion of the ledge that hangs down.
- (E) Closeup view of hairs that were left where he rubbed.



Bobcat retromingent spraying a clumped birch

Just a fine mist – mere droplets of urine upon a suitable in-your-face object on the trail – this is but one of several ways in which a Canada lynx or bobcat delivers a powerful long-lasting scent message. “Retromingent spraying” is familiar to anyone who’s observed an un-neutered tom housecat backward spraying the barn door or car tire. Curiously, for domestic cats, lynxes and bobcats alike, the “tom cat odor” actually gets stronger with a little time – hence its communicative value. Researchers have discovered that sulphur-based amino acid called “felinine” increases its odor-causing properties with the microbial and oxidative breakdown of its chemical contents. Our Keeping Track students have joined me in sniffing a certain tom bobcat’s spray mark for nearly a month before our noses could no longer pick up the felinine scent. Rotten stumps, the undersides of elevated logs, or a cliff wall beneath a rock overhang are preferred spraying locations because the urine scent message is better absorbed, protected from the elements and delivered over time. Combine just one urine spray scent post with dozens of similar spritzes over the



course of a randy tom lynx or bobcat’s sauntering search and you will appreciate that this behavior is the mate seeking season’s *piss de resistance*!

Without a snow pack for increased track detection, the “scrape” is the most frequently found wild felid sign. Scrapes are most likely to be found along ridgelines, at intersecting game trails or in the immediate vicinity of cliff habitats where my research has demonstrated bobcats locate one another and court during the breeding season. A bobcat’s hind feet push back dirt, pine needles, sand or leaves into a slight mound, upon which the felid will urinate and sometimes defecate. A bobcat’s scrape appears as a shallow rectangular trough roughly four to 5 and 1/2 inches wide. The scrape’s scuffed materials culminate in a small pile. This is both visually conspicuous and olfactorily useful as the materials both absorb and elevate the scent. In addition to urine, secretions from interdigital glands on the paws add to the mark’s scent contents.

A horizontal log or upright tree stem also serves as a selected medium for the bobcat or lynx’s scent secretions exuded from interdigital glands. Picture the wild cat crouched atop a log, in position to grasp and rake it



with repeated backward pulling motions of the forelimbs. In addition to stopping the wood with its claws and thereby removing pieces of the claws’ worn keratin, the functional purpose of **claw raking** is to scent mark. No matter how intently the felid works at shredding the wood he or she is just as pre-occupied with continually sniff - checking the roughened surface – thereby verifying the presence of the scent mark. The claw shredded wood is not only visually conspicuous; its absorbent surface collects the cat’s pedal scent and posts it for other cats’ olfactory inspection.



Catherine Morris of Quebec studies fresh scrape on Bobby Knob. Note feces on scrape.



Whitetail deer use scent marking to maintain a given herd's complex social organization. Each deer's chemo-communications convey its identity, social status and dominance ranking within the herd. Researchers have concluded that semiochemicals posted on rubs and scrapes serve to "prime" and valuably synchronize the onset of estrus in does. Other scent signals are believed to suppress the reproductive urges of subordinate bucks.



Shredded wood embedded in antler pearlulations

Researchers Larry Marchington, Karl Miller and colleagues of the University of Georgia documented that a mature whitetail buck makes an average of 2 to 3 rubs per day for a period of four months. Knobby pearlulations at the bases of his antlers combine with sharp pointed antlers and powerful neck muscles to reduce a 4 inch diameter ash trunk to shreds. Brightly colored exposed xylem and abraded wood and bark surfaces call attention to the rub; but the real purpose is to prompt olfactory inspection of the scent messages which are posted there. While making a rub a buck will repeatedly press his nose to the tree and sniff the surface he has debarked and scored with his antlers and rubbed with his forehead. His forehead's skin glands smear his unique chemical signature on the tree. Ash, cherry and willow species as well as balsam fir, elderberry and speckled alder are among the preferred rub species in the northeast.



This doe (*left*) is in the process of tasting the scent left by the buck on his rub. Non-volatile scent molecules can then be transferred by her tongue to the vomeronasal organ located on the roof of her mouth. The buck's chemical signals are then pumped to the hypothalamus, a specialized part of the brain that regulates hormones and influences reproductive readiness.

This dandy buck, shown above, has just made her a rub!

Finding the Missing Moose

Story and Photos by Susan C. Morse

Twenty-five years ago, I was planting trees on a suburban property above Shelburne Point in Vermont when I suddenly came face-to-face with a skittish young bull moose. He curiously looked me over for a minute, then whirled away and trotted to the shore of Lake Champlain. He didn't hesitate. He jumped into the water and started swimming. When I lost sight of him far across the lake, he was well on his way toward the Adirondacks.

Moose populations in the Northeast were unnaturally dense back then, and have since declined substantially. Although you're less likely now to encounter a moose wandering through a suburban landscape, you can still find them in prime summer habitat. These areas offer opportunities for cooling, concealment cover and relief from insects, and most important, provide moose with access to abundant food biomass. In summer, moose consume approximately twice as much as they do in winter—up to 45 pounds of leaves, stems, buds, flowers, and other new growth. Weight gain, stored fat, mineral nutrients, and protein are “banked” for winter's period of negative energy balance.



The moose summer diet includes dozens of species of lush forbs, as well as new leaves of woody trees and shrubs. Moose, like this handsome mature cow, spend hours immersed in wetland habitats eating the submerged stems, leaves, and rhizomes of numerous aquatic plants. These plants provide an important source of microelements, including sodium, potassium, phosphorus, and calcium, as well as crude protein. Among a moose's preferred northeastern wetland forage plants are: pondweed, common marsh marigold, water shield, yellow pond lily, fragrant water lily, and arrowhead.



Moose are still shedding their winter coats in early summer, and if you look closely in areas where you suspect they've been walking, you may notice long, coarse guard hairs and fine undercoat hairs snagged on wood. In this photo, high school students inspect hairs that a mature bull left as it ducked under a snag. The students, from Guangzhou, China, were participating in GREEN ACROSS THE WORLD®, a cultural and environmental exchange program based in Vermont.



Moose calves are the fastest-growing infants of all terrestrial mammals in North America, gaining an average of 1 to 2 pounds per day during their first month, and 3 to 5 pounds thereafter. After consuming approximately a half-gallon of its mother's rich milk each day for the first 3 to 4 weeks, a calf will gradually begin to sample some tender plant foods. Just as gradually, beneficial bacterial fauna will develop within the calf's digestive system. Some of the bacterial organisms may be picked up through direct oral contact with the mother's saliva, or indirectly, as a calf ingests plant parts that the cow has partially browsed. Calves may also "seed" their gut with bacteria by eating fecal matter (*left, bottom*) deposited by the mother and possibly other moose.



A calf's Autumn weight may be between 300 and 400 pounds, with its potential weight as a healthy yearling closer to 500 to 600 pounds. In our region, cows give birth within thick, young conifer stands next to wetlands where they are feeding. The few birthing areas I have studied were situated near wetland edges, but on elevated terrain that probably enhanced the cow's ability to detect enemies. To avoid disturbing the cow and calf, no attempt should be made to look for birthing sites until well after they've been abandoned. Usually by midsummer it is safe to scout for this sign, which includes tracks, scat, and the worn-looking and flattened appearance of vegetation and ground where the birthing and nursing occurred. While on a diet of wetland plants in the summer, moose scat is loose, like dairy cows', and less inclined to form pellets.

"If you want to find the missing moose, find the food."

Find where willow and poplar species grow, and look there. New and old browse sign will be there if moose are present. The following pages illustrate the appearance of moose feeding sign in different seasons. Across North America, willow species are a nutritious mainstay for moose in spring and summer.



Left: "Scat is loose" in spring and summer. Forbs and wetland plants can cause fecal matter to look more like that of a dairy cow.



Left: Autumn and winter's woody browse results in feces that look like they're full of sawdust and are formed into larger and smaller pellets.

Right: New and old browse sign on willow species is a sure way to find the missing moose.





A, B: Flood plain habitats, such as this riparian area in Wyoming, provide hungry moose with abundant willow forage. Many willows are easily identified because their bud arrangement is “appressed” with buds lying close or flat against the stem (see *inset*).

C, D: You’re certain to find moose sign in habitats where willow species proliferate. There are dozens of individual and hybrid species of willows that grow in the Northeast, and collectively, they’re some of the moose’s favorite summer forage plants. Tender, new leaves and budding stems provide a moose with higher amounts of protein and vitamins than all other woody plants. To identify browsed twigs, look for rough-edged stems. Moose lack upper incisors and can only partially cut twigs with the sharp, pointed edges of cusps that rise above the grinding surface of each premolar tooth. The ragged appearance of the bitten portion shows that moose must sometimes crush and partially tear away the desired bud or stem.



E, F: To browse shrubs and trees with out-of-reach branches, moose will break or bend their trunks by straddling them. These toppled “walkovers” allow moose to browse the nutritious topmost tissues of maple, willow, white birch, pin cherry, balsam fir, and hemlock. To confirm that a broken tree is a “walkover,” inspect the broken branches and browsed twigs. Notice also the presence of moose hairs on these Hemlock (*E*) and Green Alder (*F*) specimens.



Early Spring in the Rocky Mountain west is still snowbound. If you want to find the moose, find the food. *Top*, two moose are browsing among extensive willow growth along Wyoming's Gardiner River. *Above left*, moose trails show that they are seeking food among abundant willows growing in a riparian flood plain. *Above center and right*, moose in western habitats strip bark from Quaking Aspen trees. Photo on the right shows new and old bark stripping scars.



Left: Old moose barking sign on aspen becomes very corrugated-looking as it ages and the tree's trunk expands. Incisor scraping scars can be seen above the snow line and to the height of a moose's reach. Where this happens, that section of the tree's bark tends to look more like an oak than an aspen.

Above: These Wyoming aspens show a history of being foraged. In addition, we see new incisor marks, showing us that this grove is still attractive to moose.

Below: This stand of aspen in Montana is full of historic moose barking sign. Aspen bark is rich in carbohydrates, proteins and minerals, and can provide food even before the leaves are formed.





“If you want to find moose in summer, look in wetlands”

Bulls and cows alike are great swimmers. They have been observed to dive nearly 18 feet underwater, turn their bodies around, and remain foraging underwater for a half-minute or more. When a moose suddenly surfaces, its expressive rolling eyes, cascades of water draining from pendulous ears and bulbous nose make for a particularly fun and photogenic experience.

In addition to meeting other caloric demands, bull moose must also fuel the production of antlers. A mature bull such as the one pictured above is able to produce a 50-pound rack in just 150 days. During a peak growth period, his antlers can lengthen by as much as three-quarters of an inch per day. Cows (*right*) enjoy nutritious forage in the wetlands as well. A mature cow that can eat well and gain weight will be better prepared to ovulate, conceive, and support next year’s healthy calves.

Moose have been seen swimming Icy Strait in Alaska, including a moose that was killed and consumed by a killer whale! In the early 1900s, moose may have reached Isle Royale by swimming across Lake Superior from Minnesota. Their major predator, the gray wolf, probably walked across the frozen lake from Ontario in the late 1940s.



Moose are great swimmers. In summer, they seek wetlands not only for abundant, nutritious food, but for relief from insects and heat.

Hidden Treasures

Story and Photographs by Susan C. Morse



Infant fawns in the “hider phase” remain motionless and hidden in herbaceous cover. During this early period, they can be vulnerable to predators, including black bear, bobcat, coyote, fisher and occasional birds of prey.

The luminous green of grasses and forbs in late spring and early summer never fails to remind me of an especially memorable moment with my grandfather, Norman K. Morse, a naturalist, landscaper and farmer. He was part-way through a first-cut haying when he stopped his tractor and came to retrieve me. “I have something really special for you to see, but you must not make a sound,” he said, as we snuck along the woodland edge beside the hayfield.

There, flat to the ground, a centerpiece in the highest grass, lay a days-old fawn. The only movement was a flaring of nostrils as the animal breathed. Its spotted coloration visually broke up its outline, further concealing the tiny deer from predators.

My grandfather had observed an anxious doe race along the edge of the meadow as he neared where the hidden fawn was sequestered. A wise doe will even place her newborn twins in separate bed sites - as much as 100 yards apart - to preclude the loss of both should a predator discover one. The doe’s only contact with her fawns is to nurse and groom them two to three times each day. Nutritious and digestible forbs nearby (including alfalfa and clover farm crops) assure adequate milk and the fawns’ necessary and rapid growth. Weighing 6 to 10 pounds at birth, a fawn may triple its weight in a month, and, amazingly, is able to outrun predators in even less time.

By early July, the observant tracker will find the minuscule tracks and scats of some fawns. Both will be less than half the size of adult deer’s. Rarely, we find evidence of predation.



Infant fawn track.


Once, while afield with some Vermont Youth Corps volunteers, we had quietly paused for lunch on a breezy softwood bench overlooking a beaver pond. Suddenly, a heart-wrenching bleating and bawling arose from the grassy cover. Then it abruptly ceased, presumably as the predator killed the fawn.

The next day, I inspected the area and found tracks in the nearby mud and the remains of a carcass. A bear had successfully secured a meal. Sometimes, trackers can discern patterns in tall vegetation such as trampled pathways, demonstrating a bear's zigging and zagging, back-and-forth deliberate searching for

hiding fawns. As diligently as the doe conceals her fawn, some bears have learned that a deliberate search often yields much-needed, protein-rich nutrition. For a sow who has young of her own, this is nature's plan.



1. The doe will nurse and care for her hidden treasure two or three times daily
2. hider fawn in the woods
3. hider fawn in a meadow;
4. young fawn up and foraging on jewelweed – it is now strong enough to follow its mother and outrun predators.



Tables of Contents



Wildlife & Habitats

TABLE OF CONTENTS

01	Tracking 101: Thoughts About What We Look For	355	Foxes - Gray and Red
127	The World Around Us: Setting the Stage for Wildlife	377	Moose
183	Beaver	419	Porcupine
209	Birds	429	River Otter
235	Black Bear	443	Squirrels - Gray and Red
295	Bobcat	465	Whitetail Deer
311	Canada Lynx	497	Wolf, Cougar, Coyote and Other Animals:
335	Fisher		Wolf
			Cougar
			Coyote
			American Toad
			Weasel
			Marten
			Mink
			Muskrat
			Chipmunk
			Opossum
			Woodchuck
			Raccoon
			Snowshoe Hare
			Skunk
			Pronghorn



TRACKING 101: THOUGHTS ABOUT WHAT WE LOOK FOR

02	Canid versus Felid
03	When Tracks Shine
05	Making Sense of Scent Marking
08	Claw Raking: Scratching the Surface of Wild Cats' Communications Using Interdigital Scent
14	A Scent Message from an Arizona Jaguar
19	Scratch Back
21	Raised-Leg Urination
23	Hair! Hair!
27	What's the Scoop on Poop?
30	Clods, Wedgies and Imprints – Deer and moose
33	Summers's Other Nest Builders
36	Spring Cleaning
39	A Claw Comparison: How Claw Shape Relates to the Livelihoods of Bears – Black, brown, and polar bear
42	When a Wild Cat Kills a Deer PG Parental Guidance Advised
49	Forest Forensics PG Parental Guidance Advised



THE WORLD AROUND US - SETTING THE STAGE FOR WILDLIFE

- 128 Soft Serve: Autumn's Unheralded Mast Species
- 134 A Seed's Promise
- 144 High-Hanging Fruit: Boom and Bust Seed Crops of Conifers
- 153 Cavities are Good
- 162 Stump Sitters
- 171 Mushrooms on the Menu
- 179 Sign of Wee Beasties in Spring



BEAVER

- 184 Beavers at Work
- 190 Beavers at Home for the Winter
- 197 Busy Beavers



BIRDS

- 210 Fun in the North Woods with Canada Jays
- 217 Raven Partnerships
- 222 Timberdoodle
- 225 A Different Drummer
- 228 Snow Angels
- 230 Snow Birds: Staying North for the Winter



BLACK BEAR

236	Black Bear Mark Trees	263	Bear Bites and Birches
239	Grazing Bears	265	Bear Dens
243	Bears in Spring Wetlands	268	Hungry as a Bear? Try Browse and Salad
246	Babysitter Trees	270	Bear Tracks on Trees
248	Mast Miracles	272	Bear Families in Spring
251	Black Bear Day Beds	276	The Ap-Peel of Cambium
254	Grubbing	278	Looking Closely at Bear Nests
258	Bear Nests	281	Black Bear Refuge Trees and Babysitter Trees
260	Bear with Us: Signs of Black Bears too Close to Home	286	Curious Sign
		291	Reaching Back for Answers



BOBCAT

- 296 Bedside Stories
- 298 Bobcat Scrapes
- 301 Cat Communique
- 303 Bobcat Betrothals
- 305 Bobcats Searching for Mates



FISHER

- 336 Fisher Forays
- 338 A Fisher's Stash
- 340 Fishers Aren't Cats
- 342 True Grit
- 346 Getting on the Stick
- 350 Lessons from a Fisher



FOXES - GRAY AND RED

- 356 A Fox Meets its Match
- 358 New Life on the Edge
- 360 Wee Beasties Watch Out!
- 364 Is it Gray Fox or Red?
- 368 Not all Foxes are Red
- 371 Red Fox Caching



Gray fox



Red fox



© Susan C. Morse

Red fox



Reproduction Prohibited
Gray fox and pup

MOOSE

378	Spring is When the Moose Come Down	393	Declining Moose Populations: What does the Future Hold?
382	Wallowing in Love	401	Relief from Herbivory
384	Do Moose Bark?	405	Moose Rub Announcements
386	Twig Eater	408	Cow Moose Rubs on Trees
388	Blood in the Snow	412	Finding the Missing Moose
390	Winter Ticks Take Their Toll		



PORCUPINE

420 Porcupine Signs of Spring

423 Shed Hunters

425 Nip Twigs



RIVER OTTER

- 430 Feet Make Tracks
- 432 Fisher or Otter?
- 434 Raccoon or Otter?
- 436 Twist and Spraint
- 440 Slippery Business, or is it Pleasure?



SQUIRRELS - GRAY AND RED

- 444 Sap Taps
- 446 Red Squirrel's Stashes, Cashes and Trash
- 453 Gray Squirrel Caching and Feeding Sign
- 458 Putting Food Up
- 460 Manna from Heaven



Fox squirrel *Sciurus niger*



© Susan C. Morse

Red squirrel

Reproduction Prohibited
Gray squirrel

WHITETAIL DEER

466 Sorting out the Subtleties of the Rut
468 Velvet “Rub Out”
473 What’s the Rub?
476 Hidden Treasures
478 Breakfast in Bed

480 Hardship in the Deeryard
483 Marrow Core Analysis
486 Getting to the Bottom of the Scrape
492 Signposts – What’s the Rub?



WOLF, COUGAR, COYOTE AND OTHER ANIMALS

498	When the Wolf Returns	571	Chipmunk Surprises
513	Cougar Conundrum	573	Virginia Opossum
542	Omnivorous Coyote	576	The Intriguing Woodchuck
549	Keeping Track of American Toads	578	Raccoon Ramblings
552	Weasel: White on White	581	Snowshoe Hare
554	Weasels Down Under	583	Snowshoe Hares at the Edge of Spring
556	A Marten Banquet	589	Skunked!
560	Mink: Water Weasel	591	Skunk: Nose Probes and Digs
564	Mink: Summer on the Beach	593	Pronghorn Antelope – An American Original
567	The Resourceful Muskrat		






Keeping Track®

P. O. Box 1115
Richmond, VT 05477
(802)434-7000

www.keepingtrack.org
info@keepingtrack.org

(c) Susan C. Moore. All rights reserved. Reproduction prohibited

Reserve the Complete Volume Today



Pre-Order Here